

Table 1.22C Received Mental Health Treatment/Counseling and/or Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse Status and Past Year Level of Mental Illness: Standard Errors of Numbers in Thousands, 2008 and 2009

Illicit Drug or Alcohol Dependence or Abuse Status/Level of Mental Illness	Mental Health Treatment or Substance Use Treatment at a Specialty Facility ^{1,2}	Mental Health Treatment or Substance Use Treatment at a Specialty Facility ^{1,2}	Mental Health Treatment Only ¹	Mental Health Treatment Only ¹	Substance Use Treatment at a Specialty Facility Only ²	Substance Use Treatment at a Specialty Facility Only ²	Mental Health Treatment and Substance Use Treatment at a Specialty Facility ^{1,2}	Mental Health Treatment and Substance Use Treatment at a Specialty Facility ^{1,2}
	(2008)	(2009)	(2008)	(2009)	(2008)	(2009)	(2008)	(2009)
DEPENDENCE OR ABUSE	246	249	208	215	83	79	94	91
Any Mental Illness ³	239	195	195	175	119	55	84	72
Serious Mental Illness	105	124	91	111	28	16	51	56
Moderate Mental Illness ³	108	88	99	81	32	18	21	28
Mild Mental Illness ³	162	120	107	104	105	50	57	40
No Mental Illness ³	240	144	229	120	57	54	46	54
NO DEPENDENCE OR ABUSE	679	635	674	627	75	90	49	74
Any Mental Illness ³	664	420	660	413	79	38	50	63
Serious Mental Illness	236	261	231	259	16	10	36	31
Moderate Mental Illness ³	255	196	252	190	10	33	18	18
Mild Mental Illness ³	470	280	465	274	71	16	29	51
No Mental Illness ³	689	449	685	444	94	73	20	40

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: Mental Health Findings*.

¹ Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

² Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use.

³ In 2008, a split-sample design assigned adults aged 18 or older randomly to one of two impairment scales, the World Health Organization Disability Assessment Schedule (WHODAS) or the Sheehan Disability Scale (SDS). For comparability purposes, estimates for Any Mental Illness, Moderate Mental Illness, Mild Mental Illness, and No Mental Illness for 2008 are based only on the WHODAS half-sample. All estimates for 2009, as well as 2008 estimates for Serious Mental Illness, are based on the full sample. For details, see Section B.4.3 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008 and 2009.

Table 1.22D Received Mental Health Treatment/Counseling and/or Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse Status and Past Year Level of Mental Illness: Standard Errors of Percentages, 2008 and 2009

Illicit Drug or Alcohol Dependence or Abuse Status/Level of Mental Illness	Mental Health Treatment or Substance Use Treatment at a Specialty Facility ^{1,2} (2008)	Mental Health Treatment or Substance Use Treatment at a Specialty Facility ^{1,2} (2009)	Mental Health Treatment Only ¹ (2008)	Mental Health Treatment Only ¹ (2009)	Substance Use Treatment at a Specialty Facility Only ² (2008)	Substance Use Treatment at a Specialty Facility Only ² (2009)	Mental Health Treatment and Substance Use Treatment at a Specialty Facility ^{1,2} (2008)	Mental Health Treatment and Substance Use Treatment at a Specialty Facility ^{1,2} (2009)
	DEPENDENCE OR ABUSE	1.02	1.00	0.91	0.91	0.40	0.38	0.45
Any Mental Illness ³	2.25	1.55	2.09	1.52	1.43	0.60	1.04	0.80
Serious Mental Illness	2.42	2.48	2.70	2.70	1.12	0.55	1.90	1.86
Moderate Mental Illness ³	4.45	3.36	4.38	3.32	1.75	0.93	1.14	1.44
Mild Mental Illness ³	3.43	2.25	2.57	2.14	2.58	1.15	1.44	0.94
No Mental Illness ³	1.88	1.13	1.85	0.97	0.50	0.45	0.40	0.45
NO DEPENDENCE OR ABUSE	0.30	0.28	0.30	0.28	0.04	0.04	0.02	0.04
Any Mental Illness ³	1.40	0.88	1.40	0.86	0.22	0.11	0.14	0.17
Serious Mental Illness	1.77	1.85	1.78	1.88	0.22	0.12	0.48	0.37
Moderate Mental Illness ³	2.86	1.99	2.83	1.95	0.14	0.46	0.25	0.26
Mild Mental Illness ³	1.83	1.13	1.82	1.11	0.33	0.08	0.14	0.24
No Mental Illness ³	0.38	0.25	0.38	0.25	0.06	0.04	0.01	0.02

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: Mental Health Findings*.

¹ Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

² Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use.

³ In 2008, a split-sample design assigned adults aged 18 or older randomly to one of two impairment scales, the World Health Organization Disability Assessment Schedule (WHODAS) or the Sheehan Disability Scale (SDS). For comparability purposes, estimates for Any Mental Illness, Moderate Mental Illness, Mild Mental Illness, and No Mental Illness for 2008 are based only on the WHODAS half-sample. All estimates for 2009, as well as 2008 estimates for Serious Mental Illness, are based on the full sample. For details, see Section B.4.3 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008 and 2009.

Table 1.22P Received Mental Health Treatment/Counseling and/or Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 18 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse Status and Past Year Level of Mental Illness: P Values from Tests of Differences of Percentages, 2009 Versus 2008

Illicit Drug or Alcohol Dependence or Abuse Status/Level of Mental Illness	Mental Health Treatment or Substance Use Treatment at a Specialty Facility ^{1,2} (2008)	Mental Health Treatment or Substance Use Treatment at a Specialty Facility ^{1,2} (2009)	Mental Health Treatment Only ¹ (2008)	Mental Health Treatment Only ¹ (2009)	Substance Use Treatment at a Specialty Facility Only ² (2008)	Substance Use Treatment at a Specialty Facility Only ² (2009)	Mental Health Treatment and Substance Use Treatment at a Specialty Facility ^{1,2} (2008)	Mental Health Treatment and Substance Use Treatment at a Specialty Facility ^{1,2} (2009)
	DEPENDENCE OR ABUSE	0.2049		0.3657		0.8039		0.4513
Any Mental Illness ³	0.5562		0.8776		0.6205		0.1397	
Serious Mental Illness	0.5844		0.6044		0.1016		0.4487	
Moderate Mental Illness ³	0.8727		0.7494		0.8871		0.5322	
Mild Mental Illness ³	0.3884		0.4872		0.8970		0.3741	
No Mental Illness ³	0.5740		0.9267		0.5032		0.4638	
NO DEPENDENCE OR ABUSE	0.6109		0.4733		0.5359		0.2874	
Any Mental Illness ³	0.9262		0.9508		0.5318		0.3661	
Serious Mental Illness	0.8429		0.5955		0.5042		0.2660	
Moderate Mental Illness ³	0.5911		0.8279		0.1100		0.3395	
Mild Mental Illness ³	0.7896		0.7879		0.3218		0.2225	
No Mental Illness ³	0.2537		0.2195		0.9428		0.0963	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: Mental Health Findings*.

¹ Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

² Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use.

³ In 2008, a split-sample design assigned adults aged 18 or older randomly to one of two impairment scales, the World Health Organization Disability Assessment Schedule (WHODAS) or the Sheehan Disability Scale (SDS). For comparability purposes, estimates for Any Mental Illness, Moderate Mental Illness, Mild Mental Illness, and No Mental Illness for 2008 are based only on the WHODAS half-sample. All estimates for 2009, as well as 2008 estimates for Serious Mental Illness, are based on the full sample. For details, see Section B.4.3 in Appendix B of the *Results from the 2009 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008 and 2009.